

# Graph $C^*$ -Algebras with Real Rank Zero

Ja A. Jeong<sup>1</sup>

BK 21, Mathematical Sciences Division, Seoul National University, Seoul 151–742, Korea

E-mail: [jaeong@math.snu.ac.kr](mailto:jaeong@math.snu.ac.kr)

and

Gi Hyun Park<sup>2</sup>

Department of Mathematics, Hanshin University, KyungKiDo, 447–792, Korea

E-mail: [ghpark@hanshin.ac.kr](mailto:ghpark@hanshin.ac.kr)

Communicated by D. Sarason

View metadata, citation and similar papers at [core.ac.uk](http://core.ac.uk)

Given a row-finite directed graph  $E$ , a universal  $C^*$ -algebra  $C^*(E)$  generated by a family of partial isometries and projections subject to the relations determined by  $E$  is associated to the graph  $E$ . The Cuntz–Krieger algebras are those graph  $C^*$ -algebras associated to some finite graphs. We prove that a graph  $C^*$ -algebra  $C^*(E)$  has real rank zero in the sense that the set of invertible self-adjoint elements is dense in the set of all self-adjoint elements in  $C^*(E)$  (or in the unitization of  $C^*(E)$  if  $C^*(E)$  is nonunital) if and only if  $E$  satisfies a loop condition (K) that is analogous to the condition for a finite  $\{0, 1\}$  matrix  $A$  under which Cuntz analyzed the ideal structure of the Cuntz–Krieger algebra  $\mathcal{O}_A$ . © 2002 Elsevier Science (USA)

*Key Words:* real rank; directed graph; graph  $C^*$ -algebra.

## 1. INTRODUCTION

For a  $\{0, 1\}$ -matrix  $A$  satisfying a certain condition (I), Cuntz and Krieger [3] construct the Cuntz–Krieger algebra  $\mathcal{O}_A$  and Cuntz [2] determines the ideal structure of  $\mathcal{O}_A$  for a matrix  $A$  satisfying condition (II) which is stronger than (I). Graph  $C^*$ -algebras [13] for locally finite directed graphs with no sinks are introduced as an extended version of the Cuntz–Krieger algebras. In [12] Kumjian *et al.* define the graph

<sup>1</sup> Supported by BK21, KRF 2000-015-DP0019, and KOSEF 2000-0-101-001-2.

<sup>2</sup> This paper was accomplished with research fund provided by Korean Council for University Education, support for 2000 Domestic Faculty Exchange.

$C^*$ -algebra  $C^*(E)$  for any countable row-finite directed graph  $E$  possibly with sinks and prove that if  $E$  satisfies a certain loop condition (L), which is analogous to the condition (I), then an analogue of the Cuntz–Krieger uniqueness theorem holds for  $C^*(E)$ . A graph  $C^*$ -algebra  $C^*(E)$  is generated by a Cuntz–Krieger  $E$ -family of projections and partial isometries subject to the relations determined by  $E$  or by the edge matrix of  $E$  if  $E$  has no sinks. Moreover  $C^*(E)$  is a universal  $C^*$ -algebra for Cuntz–Krieger  $E$ -family. In [13], a groupoid  $\mathcal{G}$  associated to a locally finite directed graph  $E$  with no sinks is considered and it is shown in the same paper that the groupoid  $C^*$ -algebra  $C^*(\mathcal{G})$  is a universal  $C^*$ -algebra generated by a Cuntz–Krieger  $A$ -family of projections and partial isometries, where  $A$  is the edge matrix for the graph  $E$ . It then follows that the two  $C^*$ -algebras  $C^*(\mathcal{G})$  and  $C^*(E)$  are isomorphic, and hence some knowledge for groupoid  $C^*$ -algebras could be applied to analyze the ideal structure of graph  $C^*$ -algebras when  $E$  satisfies condition (K), which is analogous to the condition (II) of a matrix  $A$  generating a Cuntz–Krieger algebra  $\mathcal{O}_A$ .

Since graph  $C^*$ -algebras  $C^*(E)$  have partial isometries and projections as their generators one might expect that most of the graph  $C^*$ -algebras would have enough projections and they might have real rank zero. In fact, the Cuntz algebras  $\mathcal{O}_n$  are known to have real rank zero since they are purely infinite and simple [15], but the Toeplitz algebra and its quotient algebra  $C(\mathbb{T})$  are graph  $C^*$ -algebras with real rank greater than 0. Thus the question in view of abundance of projections is to find a condition in terms of the graph under which the associated graph  $C^*$ -algebra has real rank 0. It is proved in [7] that if  $C^*(E)$  has real rank zero then  $E$  should satisfy condition (K), and the converse is true if  $C^*(E)$  has finitely many ideals. In this paper we show that for a locally finite directed graph  $E$  the graph  $C^*$ -algebra  $C^*(E)$  has real rank zero exactly when  $E$  satisfies condition (K). Given a directed graph  $E$ , it is immediate to see whether  $E$  satisfies condition (K). Hence our theorem provides a very convenient way to know that the real rank of a graph  $C^*$ -algebra will be zero or not.

It is known [8] that a separable nuclear simple  $C^*$ -algebra  $A$  is purely infinite if and only if  $A \simeq A \otimes \mathcal{O}_\infty$ . It then follows from our theorem and a result in [6] that if a graph  $C^*$ -algebra  $C^*(E)$  is purely infinite in the sense of [9] (simple or nonsimple) its real rank is zero. Hence we have that  $C^*(E)$  is purely infinite if and only if  $C^*(E) \simeq C^*(E) \otimes \mathcal{O}_\infty$  by [10, Corollary 9.4].

## 2. PRELIMINARIES

A directed graph consists of vertices and directed edges between them. More formally, a directed graph  $E$  is a quadruple  $(E^0, E^1, r, s)$ , where  $E^0$

and  $E^1$  are countable sets of *vertices* and *edges*, and  $r, s : E^1 \rightarrow E^0$  are the *range* and *source* maps, respectively.

A vertex  $v$  that emits no edges is called a *sink*. A graph  $E$  is *row-finite* if every vertex emits only finitely many edges. A row-finite graph  $E$  is *locally finite* if every vertex receives only finitely many edges and *finite* if  $E^0$  is a finite set.

We call a sequence  $\alpha = \alpha_1 \alpha_2 \cdots \alpha_n$  of edges  $\alpha_i$ , where  $r(\alpha_i) = s(\alpha_{i+1})$  for  $1 \leq i \leq n-1$ , a *path* of length  $|\alpha| = n$ . A finite path  $\alpha$  with  $|\alpha| > 0$  is called a *loop* based at  $v$  if  $s(\alpha) = r(\alpha) = v$ . We denote  $\alpha^0$  the set of vertices on the loop  $\alpha = \alpha_1 \alpha_2 \cdots \alpha_n$ , that is,  $\alpha^0 = \{v \in E^0 \mid v = s(\alpha_i), 1 \leq i \leq n\}$  as in [6]. A loop  $\alpha$  is *simple* if the vertices  $s(\alpha_i)$ ,  $1 \leq i \leq |\alpha|$ , are distinct. For a subgraph  $F$  of  $E$ , an *exit* of  $F$  is an edge  $e$  of  $E$  with  $s(e) \in F^0$  and  $e \notin F^1$ .

Let  $E^n$  be the set of paths of length  $n$ . By convention  $E^0$  is identified with the set of paths of length 0. The range map  $r$  and the source map  $s$  extend to  $E^* = \bigcup_{n \geq 0} E^n$  by  $r(\alpha) = r(\alpha_{|\alpha|})$  and  $s(\alpha) = s(\alpha_1)$ .

A graph  $E$  is said to satisfy *condition (L)* if every simple loop in  $E$  has an exit, and *condition (K)* if for each vertex  $v$  on a loop there exist at least two distinct loops  $\alpha, \beta$  based at  $v$  such that  $r(\alpha) = r(\beta) = s(\alpha) = s(\beta) = v$ ,  $r(\alpha_i) \neq v$  for  $1 \leq i < |\alpha|$  and  $r(\beta_j) \neq v$  for  $1 \leq j < |\beta|$ .

Given a row-finite graph  $E$ , a *Cuntz–Krieger  $E$ -family* consists of a family of mutually orthogonal projections  $\{p_v : v \in E^0\}$  indexed by vertices and a family of partial isometries  $\{s_e : e \in E^1\}$  indexed by edges satisfying the *Cuntz–Krieger  $E$ -relations*:

$$s_e^* s_e = p_{r(e)} \quad \text{for } e \in E^1,$$

$$p_v = \sum_{\{e : s(e) = v\}} s_e s_e^* \quad \text{for } v \in E^0, \quad \text{which is not a sink.}$$

In [12] Kumjian *et al.* showed that there exists a  $C^*$ -algebra  $C^*(E)$  generated by a Cuntz–Krieger  $E$ -family  $\{s_e, p_v\}$ , which is universal in the sense that for any Cuntz–Krieger  $E$ -family  $\{t_e, q_v\}$  in a  $C^*$ -algebra  $A$ , there exists a  $*$ -homomorphism  $\pi : C^*(E) \rightarrow A$  such that  $\pi(s_e) = t_e$  and  $\pi(p_v) = q_v$ . This  $C^*$ -algebra  $C^*(E)$  is called the *graph  $C^*$ -algebra of  $E$* . It is useful to note that the linear span of the set  $\{s_\alpha s_\beta^* \mid \alpha, \beta \in E^*, r(\alpha) = r(\beta)\}$  is dense in  $C^*(E)$  and the finite sums of projections  $\{p_v\}$  forms an approximate unit for  $C^*(E)$ . Moreover graph  $C^*$ -algebras are nuclear since they are stably isomorphic to the crossed products of AF algebras by the integer group  $\mathbb{Z}$  (see [11]).

### 3. IDEAL STRUCTURE OF GRAPH $C^*$ -ALGEBRAS

We need to review the ideal theory of a graph  $C^*$ -algebra  $C^*(E)$  for a directed graph  $E$  with no sinks [13].

For vertices  $v, w$  of  $E$ , we write  $v \geq_E w$  if  $v = w$  or there is a path  $\alpha$  with  $s(\alpha) = v$  and  $r(\alpha) = w$ . Then a subset  $H$  of  $E^0$  is called *hereditary* if

$$v \geq_E w \quad \text{and} \quad v \in H \Rightarrow w \in H,$$

and *saturated* if

$$s^{-1}(v) \neq \emptyset \quad \text{and} \quad r(s^{-1}(v)) \subset H \Rightarrow v \in H.$$

Given set inclusion as the order relation, the set of all hereditary saturated subsets has the complete lattice structure. For a subset  $N$  of  $E^0$  there exists a smallest hereditary saturated subset of  $E^0$  containing  $N$  which we call the *hereditary saturation* of  $N$  and write  $\langle N \rangle$ .

If  $E$  is row-finite, every hereditary saturated subset  $H$  of  $E^0$  gives rise to a closed two-sided ideal  $I(H)$  of  $C^*(E)$ , where

$$I(H) = \overline{\text{span}}\{s_\alpha s_\beta^* \mid \alpha, \beta \in E^*, r(\alpha) = r(\beta) \in H\}.$$

Furthermore the following is known. By an ideal we always mean a closed two-sided one.

**THEOREM 3.1** ([13, THEOREM 6.6] OR [14, THEOREM 2.2]). *Let  $E$  be a locally finite directed graph with no sinks. Then the map  $H \mapsto I(H)$  described above is injective, and the quotient algebra  $C^*(E)/I(H)$  is isomorphic to  $C^*(F)$  of the directed graph  $F := (E^0 \setminus H, \{e \mid r(e) \notin H\})$ . The ideal  $I(H)$  is strong Morita equivalent to  $C^*(G)$  of the directed graph  $G := (H, \{e \mid s(e) \in H\})$ . Moreover, if  $E$  satisfies condition (K) then the map  $H \mapsto I(H)$  is an isomorphism of the lattice of hereditary saturated subsets of  $E^0$  onto the lattice of ideals in  $C^*(E)$ .*

Using this theorem we will see the ideal structure of graph C\*-algebra  $s$  by investigating hereditary saturated subsets of vertices of the graphs. In the following we define some notions on graphs which will facilitate counting the number of hereditary subsets of vertices in a graph.

**DEFINITION 3.1.** Let  $F$  be a subgraph of  $E$ . The *loop completion*  $\ell_E(F)$  of  $F$  in  $E$  is a subgraph of  $E$  that is the union of  $F$  and all the loops of  $E$  based at the vertices of  $F$ . The *loop contraction*  $\ell_c(F)$  of  $F$  is a graph obtained by shrinking each loop in  $F$  to one consisting of only one edge.

More formally, the loop contraction can be defined as follows. First we define relations  $\sim_0$  and  $\sim_1$  on vertex set and edge set of  $F$ , respectively;

$$\begin{aligned} v \sim_0 w &\Leftrightarrow \ell_F(\{v\}) = \ell_F(\{w\}), \\ e \sim_1 f &\Leftrightarrow s(e) \sim_0 s(f), r(e) \sim_0 r(f). \end{aligned}$$

These are easily seen to be equivalence relations. We extend the range map and the source map to those on the equivalence classes naturally;  $r([e]) = [r(e)]$  and  $s([e]) = [s(e)]$ . Then the quadruple  $[F] := (F^0 / \sim_0, F^1 / \sim_1, r, s)$  becomes a directed graph and has, if any, only simple loops consisting of one edge. Then the loop contraction  $\ell c(F)$  of  $F$  is defined to be the graph  $[F]$ .

Note that under the quotient maps  $f_0: v \in F^0 \mapsto [v] \in F^0 / \sim_0$  and  $f_1: e \in F^1 \mapsto [e] \in F^1 / \sim_1$ , the order relation of vertices in  $F$  is preserved:

$$v \geq_F w \Leftrightarrow [v] \geq_{\ell c(F)} [w].$$

Thus  $F$  and its loop contraction  $\ell c(F)$  have the isomorphic lattice structure of hereditary subsets of vertices.

**LEMMA 3.1.** *Let  $E$  be a locally finite directed graph and  $F$  a subgraph of  $E$ .*

1. *If  $F$  is a finite subgraph of  $E$ , then the number of hereditary subsets of  $\ell_E(F)^0$  is finite.*

2. *Let  $G$  be a graph obtained from  $F$  by adding some edges between some pairs of vertices of  $F$ . Then the number of hereditary subsets of  $G^0$  is less than or equal to that of  $F^0$ .*

*Proof.* (1) Since the cardinality of the vertex set  $\ell c(\ell_E(F))^0$  is not larger than that of  $F^0$  and two graphs  $\ell c(\ell_E(F))$  and  $\ell_E(F)$  have the isomorphic lattice structure of hereditary vertex subsets, the number of hereditary subsets of  $\ell_E(F)^0$  is finite.

(2) Adding edges implies that

$$v \geq_F w \Rightarrow v \geq_G w.$$

Let  $H$  be a  $G$ -hereditary subset, that is, a hereditary subset of  $G^0$ . Then

$$\begin{aligned} v \geq_F w, \quad v \in H &\Rightarrow v \geq_G w, \quad v \in H \\ &\Rightarrow w \in H. \end{aligned}$$

Therefore every  $G$ -hereditary subset is a  $F$ -hereditary subset. ■

**DEFINITION 3.2.** Let  $F$  be a subgraph of  $E$ . The *exit completion*  $F_e$  of  $F$  is a subgraph of  $E$  obtained by adding to  $F$  all the edges  $e \in E^1$  such that  $s(e) = s(f)$  for some  $f \in F^1$  and their range vertices  $r(e)$ . If  $F = F_e$  we call  $F$  *exit complete*.

Note that  $F_e$  does not necessarily contain all the exits of  $F$ .

**PROPOSITION 3.1.** *Let  $F$  be an exit complete subgraph of  $E$  and  $B$  be the  $C^*$ -subalgebra of  $C^*(E)$  generated by  $\{s_e, p_v \mid e \in F^1, v \in F^0\}$ . Then there is a  $*$ -isomorphism  $h: C^*(F) \rightarrow B$  such that*

$$h(t_e) = s_e, \quad h(q_v) = p_v, \quad e \in F^1, \quad v \in F^0,$$

where  $\{t_e \mid e \in F^1\} \cup \{q_v \mid v \in F^0\}$  is the Cuntz–Krieger  $F$ -family generating  $C^*(F)$ .

*Proof.* Let  $\{s_e \mid e \in E^1\} \cup \{p_v \mid v \in E^0\}$  be the Cuntz–Krieger  $E$ -family generating  $C^*(E)$ . If  $v \in F^0$  then either  $v$  is a sink of  $F$  or  $s^{-1}(v) = \{e \in E^1 \mid s(e) = v\} \subset F^1$  since  $F$  is exit complete. For each nonsink  $v \in F^0$ , the Cuntz–Krieger  $E$ -relation at  $v$ ,  $p_v = \sum_{\{e \in E^1 \mid s(e) = v\}} s_e s_e^*$ , coincides with the Cuntz–Krieger  $F$ -relation  $p_v = \sum_{\{e \in F^1 \mid s(e) = v\}} s_e s_e^*$  at  $v$ . Thus the elements  $\{s_e \mid e \in F^1\} \cup \{p_v \mid v \in F^0\}$  in  $C^*(E)$  forms a Cuntz–Krieger  $F$ -family. Now let  $C^*(F)$  be represented faithfully in a Hilbert space  $H = \bigoplus_{v \in F^0} H_v$ , where  $H_v = q_v H$  is an infinite-dimensional Hilbert space corresponding to each vertex  $v \in F^0$  such that  $H_v = \bigoplus_{e \in E^1} H_e$  and  $t_e (e \in F^1)$  is an isometry of  $H_{r(e)}$  onto  $H_e$ . Similarly, for each  $w \in E^0 \setminus F^0$ ,  $f \in E^1 \setminus F^1$  take infinite dimensional Hilbert spaces  $H_f$  and  $H_w = \bigoplus_{s(f)=w} H_f$ . Put  $K = \bigoplus_{v \in E^0} H_v$ . Choose an isometry  $S_f$  of  $H_{r(f)}$  onto  $H_f$  and let  $P_w$  be the orthogonal projection from  $K$  onto  $H_w$ . Then the projections  $\{q_v, P_w \mid v \in F^0, w \in E^0 \setminus F^0\}$  and the partial isometries  $\{t_e, S_f \mid e \in F^1, f \in E^1 \setminus F^1\}$  form a Cuntz–Krieger  $E$ -family in the Hilbert space  $K$ . Hence there exists a  $*$ -isomorphism  $g$  from  $C^*(E)$  onto the  $C^*$ -algebra generated by these projections and partial isometries such that

$$h(p_v) = \begin{cases} q_v, & v \in F^0 \\ P_v, & v \in E^0 \setminus F^0 \end{cases} \quad \text{and} \quad h(s_e) = \begin{cases} t_e, & e \in F^1 \\ S_e, & e \in E^1 \setminus F^1. \end{cases}$$

Note here that  $C^*(F) = g(B)$  since  $g(s_e) = t_e$ ,  $e \in F^1$  and  $g(p_v) = q_v$ ,  $v \in F^0$ . From the universal property of the  $C^*$ -algebra  $C^*(F)$  it follows that  $B$  has the same property. Then the uniqueness of such a universal  $C^*$ -algebra implies that  $g$  should be an isomorphism from  $B$  onto  $C^*(F)$  and  $h = g^{-1}$  is the desired isomorphism. ■

#### 4. REAL RANK OF GRAPH $C^*$ -ALGEBRAS

The *real rank* of a unital  $C^*$ -algebra  $A$ , denoted by  $\text{RR}(A)$ , is defined by Brown and Pedersen [1] to be the smallest integer  $n$  such that for each  $(n+1)$ -tuple  $(x_1, \dots, x_{n+1})$  of self-adjoint elements in  $A$  and every  $\varepsilon > 0$ ,

there is an  $(n+1)$ -tuple  $(y_1, \dots, y_{n+1})$  of self-adjoint elements in  $A$  such that  $\sum y_k^2$  is invertible and

$$\left\| \sum_{k=1}^{n+1} (x_k - y_k)^2 \right\| < \varepsilon.$$

For a nonunital  $C^*$ -algebra  $A$ , its real rank is defined to be  $\text{RR}(A^\sim)$ , where  $A^\sim$  is the unitization of  $A$ .

In particular unital  $C^*$ -algebras of real rank zero are those in which every self-adjoint element can be arbitrarily closely approximated by invertible self-adjoint elements, or equivalently every hereditary  $C^*$ -subalgebra of  $A$  contains an approximate unit consisting of projections [1]. Hence  $\text{RR}(A) = 0$  implies that  $A$  contains fairly many projections. All AF algebras and all von Neumann algebras have real rank zero. Recall that a simple  $C^*$ -algebra is said to be *purely infinite* if each of its nonzero hereditary  $C^*$ -subalgebra has an infinite projection. It is known that a purely infinite simple  $C^*$ -algebra is of real rank zero [15]. For example, the Cuntz algebras  $\mathcal{O}_n$  ( $2 \leq n \leq \infty$ ) have real rank zero. Also some of the Cuntz–Krieger algebras  $\mathcal{O}_A$  are known to have real rank zero (see [7, Corollary 4.7]).

Since graph  $C^*$ -algebras  $C^*(E)$  have projections and partial isometries as generators like Cuntz–Krieger algebras one might expect that most of the graph  $C^*$ -algebras would contain enough projections and have real rank zero. The question we are interested in is to find a graph theoretic condition under which the graph  $C^*$ -algebras are of real rank zero, and the following theorem is the answer to this question.

**THEOREM 4.1.** *Let  $E$  be a locally finite directed graph. Then the following are equivalent.*

1.  $C^*(E)$  is of real rank zero.
2.  $E$  satisfies condition (K).
3.  $C^*(E)$  has no quotients containing a corner that is  $*$ -isomorphic to  $M_n(C(\mathbb{T}))$ .

A partial result has been known for some graph  $C^*$ -algebras.

**THEOREM 4.2** ([7, THEOREMS 4.3 and 4.6]). *Let  $E$  be a locally finite directed graph with no sinks. If  $\text{RR}(C^*(E)) = 0$  then  $E$  satisfies condition (K). Conversely if  $E$  satisfies condition (K) and  $C^*(E)$  has only finitely many ideals then  $\text{RR}(C^*(E)) = 0$ . In particular, if  $E$  is a finite graph then  $\text{RR}(C^*(E)) = 0$ .*

Recall that if  $E$  is a locally finite directed graph with no sinks and  $H$  is a hereditary saturated subset of  $E^0$  then the quotient algebra  $C^*(E)/I(H)$  is

isomorphic to  $C^*(F)$  where  $F := (E^0 \setminus H, \{e \mid r(e) \notin H\})$ . We simply call  $F$  a *quotient graph* of  $E$ .

**PROPOSITION 4.1.** *Let  $E$  be a locally finite graph with no sinks and  $H$  be a hereditary saturated vertex subset of  $E^0$ . If both the ideal  $I(H)$  and the quotient algebra  $C^*(E)/I(H) \cong C^*(F)$  have real rank zero, then  $RR(C^*(E)) = 0$ .*

*Proof.* By [1, Theorem 3.14], it is enough to show that every projection in the quotient algebra  $C^*(E)/I(H) \cong C^*(F)$  lifts to a projection in  $C^*(E)$ .

For a row-finite graph  $E$  with no sinks, the  $K_0$  group  $K_0(C^*(E))$  is generated, as an Abelian group, by the equivalence classes  $\{[p_v] : v \in E^0\}$  subject to the relation

$$[p_v] = \sum_{e: s(e)=v} [p_{r(e)}]$$

by [14, Theorem 3.1]. It then follows that the induced homomorphism from  $K_0(C^*(E))$  to  $K_0(C^*(E)/I(H))$  is surjective. By [1, Proposition 3.15] we see that every projection in  $C^*(E)/I(H)$  lifts to a projection in  $C^*(E)$ . ■

*Proof of Theorem 4.1.* Let  $H$  be the hereditary saturation of the sinks of  $E$ . Then the ideal  $I(H) = \overline{\text{span}}\{s_\alpha s_\beta^* \mid \alpha, \beta \in E^*, r(\alpha) = r(\beta) \in H\}$  is AF by [12, Corollary 2.2] and the quotient algebra  $C^*(E)/I(H)$  is isomorphic to  $C^*(G)$ , where  $G = (E^0 \setminus H, \{e \in E^1 \mid r(e) \notin H\})$  is a quotient graph of  $E$  with no sinks [7, Theorem 3.5]. Since  $RR(I(H)) = 0$  and every projection in  $C^*(G)$  lifts to a projection in  $C^*(E)$  by [1, Corollary 3.16] because  $K_1(I(H)) = 0$ , it follows from [1, Theorem 3.14] that  $RR(C^*(E)) = 0$  if and only if  $RR(C^*(G)) = 0$ . Also, it is easy to see that  $E$  satisfies condition (K) exactly when  $G$  does so. Thus we may assume that  $E$  has no sinks for the proof of (1)  $\Leftrightarrow$  (2).

(1)  $\Rightarrow$  (2): This follows from Theorem 4.2.

(2)  $\Rightarrow$  (1): Assume that  $E$  satisfies condition (K). Let  $s$  be a self-adjoint element of  $C^*(E)^\sim$ . Since the linear span of the set  $\{s_\alpha s_\beta^* \mid \alpha, \beta \in E^*, r(\alpha) = r(\beta)\}$  is dense in  $C^*(E)$ , there is a finite sum  $s' = \sum \lambda_{\alpha\beta} s_\alpha s_\beta^* + \lambda \cdot 1$  that approximates  $s$  close enough and is self-adjoint. Let  $F$  be the loop completion of the finite subgraph of  $E$  consisting of all the edges  $e$  of  $\alpha, \beta$  and all their vertices  $r(e), s(e)$  for  $\alpha, \beta$  appearing in the above expression of  $s'$ . Let  $F_e$  be the exit completion of  $F$ .

Now we claim that  $C^*(F_e)$  is of real rank zero. Considering the ideal  $I(\langle S \rangle)$  of  $C^*(F_e)$  generated by the sinks  $S$  of  $F_e$  as in the first paragraph of



the proof it suffices to prove that  $C^*(G)$  has real rank zero, where  $G = (F_e^0 \setminus \langle S \rangle, \{e \in F_e^1 \mid r(e) \notin \langle S \rangle\})$  is a subgraph of  $F_e$  with no sinks such that  $C^*(F_e)/I(\langle S \rangle) \simeq C^*(G)$ . Note here that the graph  $G$  satisfies condition (K) because  $G$  contains all the loops of  $F_e$  and the loops of  $F_e$  are contained in  $F$ , which satisfies condition (K) as a loop completion of some subgraph of  $E$  with (K). By Theorem 4.2 it is enough to see that  $C^*(G)$  has finitely many ideals. We will do this by showing that the number of hereditary subsets of  $G^0$  is finite.

The exit completion  $F_e$  of  $F$  is obtained by adding to  $F$  some vertices and some edges. Let  $V := F_e^0 \setminus F^0$  be the set of vertices added to  $F^0$ . Then  $V \subset \langle S \rangle$ . The set of edges in  $F_e$  added to  $F$  is the disjoint union of  $E^+ := \{e \in F_e^1 \mid r(e) \in V\}$  and  $E^\dagger := \{e \notin F^1 \mid s(e), r(e) \in F^0 \text{ and } s(e) = s(f) \text{ for some } f \in F^1\}$ . Let  $K$  be the subgraph of  $F_e$  obtained by removing the vertices in  $V$  and all edges in  $E^+$ , or equivalently that obtained by adding only the edges in  $E^\dagger$  to  $F$ . The number of hereditary subsets of  $K^0$  is finite by Lemma 3.1.

Since  $F_e^0 = G^0 \cup \langle S \rangle$ , we have  $K^0 = F_e^0 \setminus V = (G^0 \cup \langle S \rangle) \setminus V = G^0 \cup (\langle S \rangle \setminus V)$ . Now it is easy to show that for a hereditary subset  $H$  of  $G^0$ ,  $H \cup (\langle S \rangle \setminus V)$  is a hereditary subset of  $K^0$  and this correspondence is one to one between the set of hereditary subsets of  $G^0$  and the set of hereditary subsets of  $K^0$  containing  $\langle S \rangle \setminus V$ , so that the number of hereditary subsets of  $G^0$  is less than or equal to that of  $K^0$ . Therefore the number of hereditary subsets of  $G^0$  is finite.

Now let  $B$  be the  $C^*$ -subalgebra of  $C^*(E)$  generated by  $\{s_e, p_v : e \in F_e^1, v \in F_e^0\}$ . Then the element  $s'$  is in  $B^\sim$  by the choice of  $F_e$ . Since  $F_e$  is exit complete in  $E$ , by Proposition 3.1 there exists a  $C^*$ -homomorphism  $h$  from  $C^*(F_e)$  onto  $B$  such that  $h(t_e) = s_e$ ,  $h(q_v) = p_v$ , where  $\{t_e, q_v\}$  is the Cuntz–Krieger  $F_e$ -family generating the graph  $C^*$ -algebra  $C^*(F_e)$ . Let  $s''$  be an element of  $C^*(F_e)^\sim$  that can be written in the same form as  $s'$  in  $C^*(E)^\sim$ ;  $s'' = \sum \lambda_{\alpha\beta} t_\alpha^* t_\beta^* + \lambda \cdot 1 \in C^*(F_e)^\sim$ . Then there is an invertible self-adjoint element  $t$  of  $C^*(F_e)^\sim$  that approximates  $s''$  close enough since  $RR(C^*(F_e)^\sim) = 0$ . The homomorphism  $h$  maps the element  $s'' \in C^*(F_e)^\sim$  to  $s' \in B^\sim$ . Thus  $s'$  and  $h(t)$  are close enough and we conclude that the invertible self-adjoint element  $h(t)$  approximates  $s$  as close as desired. Therefore  $C^*(E)^\sim$  (and so  $C^*(E)$ ) is of real rank zero.

(1)  $\Rightarrow$  (3): Note that the property of having real rank zero passes through the quotient algebras and hereditary  $C^*$ -subalgebras, but clearly  $M_n(C(\mathbb{T}))$  does not have real rank zero.

(3)  $\Rightarrow$  (2): Suppose  $E$  does not satisfy condition (K). Then the subgraph  $G$  of  $E$  corresponding to the quotient algebra  $C^*(E)/I(H)$  clearly does not satisfy condition (K), where  $H$  is the hereditary saturation of the sinks of  $E$ .  $G$  has no sinks and there is a simple loop  $\alpha = \alpha_1 \alpha_2 \cdots \alpha_n$

such that  $\alpha$  is the only loop based at the vertex  $s(\alpha)$ . If  $\alpha$  has an exit  $e$ , the range vertex  $r(e)$  does not belong to  $\alpha^0$ . Let  $V$  be the hereditary saturation of the vertices  $\{r(e) \mid e \text{ is an exit of } \alpha\}$ . Then the quotient algebra  $C^*(G)/I(V)$  is isomorphic to a graph  $C^*$ -algebra  $C^*(F)$ , where  $F = (G^0 \setminus V, \{e \in G^1 \mid r(e) \notin V\})$ . Note that  $F$  contains the loop  $\alpha$  and  $\alpha$  has no exits in  $F$ . By [6, Lemma 4.5]  $C^*(F)$  contains a projection  $p$  such that the corner  $pC^*(F)p$  is isomorphic to  $M_n(C(\mathbb{T}))$ . If  $\alpha$  has no exits then  $G = F$  and we have the same conclusion. ■

It is known that a simple, separable, nuclear, unital  $C^*$ -algebra  $A$  is purely infinite if and only if  $A \simeq A \otimes \mathcal{O}_\infty$ , where  $\mathcal{O}_\infty$  is the Cuntz algebra generated by isometries  $\{S_n \mid n \in \mathbb{N}\}$  with orthogonal ranges (see [8], for example). For nonsimple  $C^*$ -algebras  $A$ , three notions of pure infiniteness are introduced in [9] and [10] for the purpose of characterizing nonsimple  $C^*$ -algebras absorbing the Cuntz algebra  $\mathcal{O}_\infty$  (that is,  $A \simeq A \otimes \mathcal{O}_\infty$ ). A  $C^*$ -algebra  $A$  is *purely infinite* if  $A$  has no Abelian quotients and every nonzero positive element in  $A$  is properly infinite (see [9]). It is proved in [10, Corollary 9.3] that for a separable, nuclear  $C^*$ -algebras  $A$  with an approximate unit consisting of projections  $A$  is approximately divisible and purely infinite if and only if  $A \simeq A \otimes \mathcal{O}_\infty$ . In [6], Hjelmborg proves among others that a graph  $C^*$ -algebra  $C^*(E)$  is purely infinite if and only if every vertex connects to a loop with an exit in every quotient graph  $F$ . This means that every quotient graph  $F$  satisfies condition (K) and every vertex in  $F$  connects to a loop in  $F$ . In particular we have that if  $C^*(E)$  is purely infinite then  $E$  satisfies condition (K) and hence  $C^*(E)$  has real rank zero, but for a separable, nuclear  $C^*$ -algebra  $A$  of real rank zero it is known in [10, Corollary 9.4] that  $A$  is purely infinite if and only if  $A \simeq A \otimes \mathcal{O}_\infty$ . Therefore, with the result [12, Theorem 2.4] that  $C^*(E)$  is AF if and only if  $E$  has no loops, we have the following corollary.

**COROLLARY 4.1.** *Let  $E$  be a locally finite directed graph with no sinks. Then the following are equivalent.*

1.  $C^*(E)$  is purely infinite,
2.  $C^*(E) \simeq C^*(E) \otimes \mathcal{O}_\infty$ ,
3.  $C^*(E)$  is purely infinite and approximately divisible,
4.  $C^*(E)$  has no AF quotients and  $\text{RR}(C^*(E)) = 0$ , and
5.  $C^*(E)$  has no AF quotients and  $E$  satisfies condition (K).

Note also that  $C^*(E)$  is purely infinite then every nonzero hereditary  $C^*$ -subalgebra of  $C^*(E)$  contains an infinite projection, which follows from [6, Theorem 3.1] and [12, Theorem 3.9], but this is not true for  $C^*$ -algebras that are not associated to directed graphs, in general.

## REFERENCES

1. L. G. Brown and G. K. Pedersen,  $C^*$ -algebras of real rank zero, *J. Funct. Anal.* **99** (1991), 131–149.
2. J. Cuntz, A class of  $C^*$ -algebras and topological Markov chains II: Reducible chains and the Ext-functor for  $C^*$ -algebras, *Invent. Math.* **63** (1981), 25–44.
3. J. Cuntz and W. Krieger, A class of  $C^*$ -algebras and topological Markov chains, *Invent. Math.* **56** (1980), 251–268.
4. D. Drinen, Viewing AF-algebras as graph algebras, *Proc. Amer. Math. Soc.* **128** (2000), 1991–2000.
5. D. Drinen and M. Tomforde, The  $C^*$ -algebras of arbitrary graphs, preprint, 2000.
6. J. v. B. Hjelmborg, Purely infinite and stable  $C^*$ -algebras of graphs and dynamical systems, in press.
7. J. A. Jeong, G. H. Park, and D. Y. Shin, Stable rank and real rank of graph  $C^*$ -algebras, *Pacific J. Math.* **200** (2001), 331–343.
8. E. Kirchberg, The classification of purely infinite  $C^*$ -algebras using Kasparov’s theory, *Fields Inst. Commun.*, in press.
9. E. Kirchberg and M. Rordam, Non-simple purely infinite  $C^*$ -algebras, *Amer. J. Math.* **122** (2000), 637–666.
10. E. Kirchberg and M. Rordam, Infinite non-simple  $C^*$ -algebras absorbing the Cuntz algebra  $\mathcal{O}_\infty$ , preprint, 2000.
11. A. Kumjian, Notes on  $C^*$ -algebras of graphs, in “Operator Algebras and Operator Theory,” *Comm. Amer. Math. Soc.* **228** (1998), 189–200.
12. A. Kumjian, D. Pask, and I. Raeburn, Cuntz–Krieger algebras of directed graphs, *Pacific J. Math.* **184** (1998), 161–174.
13. A. Kumjian, D. Pask, I. Raeburn, and J. Renault, Graphs, groupoids, and Cuntz–Krieger algebras, *J. Funct. Anal.* **144** (1997), 505–541.
14. D. Pask, Cuntz–Krieger algebras associated to directed graphs, in “Operator Algebras and Quantum Field Theory, Rome, July 1–6, 1996” (S. Doplicher, R. Longo, J. Roberts, and L. Zsido, Eds.), pp. 85–92, International Press, 1997.
15. S. Zhang, A property of purely infinite simple  $C^*$ -algebras, *Proc. Amer. Math. Soc.* **109** (1990), 717–720.